

239-2522

11/27/2012

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Ms Mandy Slack  
The Scotts Company  
d/b/a The Ortho Group  
PO BOX 190  
Marysville, OH 43040

**NOTIFICATION**  
NOV 27 2012

Subject: Notification for Label Revisions under PRN 98-10 and 2008-1: Environmental Hazards and other changes  
Submission date: 10/24/2012  
Product Name: Ortho Daconil 2787 Multipurpose Fungicide  
EPA Reg. No. 239-2522  
Decision Number 471797

Dear Registrant:

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2008-1 and 98-10, and finds that the label change(s) requested falls within the scope of PRN2008-1 and 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have questions concerning this letter, please call Banza Djapao at 703-305-7269 or you may call me at 703-308-9443.

Sincerely,

A handwritten signature in black ink that reads "Tony Kish".

Tony Kish  
Product Manager 22  
Fungicide Branch  
Registration Division (7504P)

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98

<b>EPA</b> United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	Identifier Number
---	--	-------------------

**Application for Pesticide - Section I**

1. Company/Product Number The Scotts Company d/b/a The Ortho Group / 239-2522	2. EPA Product Manager Tony Kish	3. Proposed Classification  <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Ortho Group / Ortho® Daconil 2787® Multipurpose Fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) The Scotts Company d/b/a The Ortho Group PO Box 190 Marysville, OH 43040  <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

**Section II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - Explain below.
--	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
 Updated the Environmental Hazards section per PR Notice 2008-1; minor labeling additions/corrections per PR Notice 98-10. See attached letter for details.  
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section III**

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container		
	If "Yes" Unit Packaging wgt.    No. per Container	If "Yes" Package wgt.    No. per Container	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
*Certification must be submitted					
3. Location of Net Contents Information		4. Size(s) Retail Container		5. Location of Label Directions	
<input type="checkbox"/> Label <input type="checkbox"/> Container				<input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		

**Section IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Mandy Slack	Title Analyst, Regulatory Affairs	Telephone No. (Include Area Code) (937) 644-7187
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received <b>(Stamped)</b>
2. Signature The Scotts Company LLC BY: <i>Mandy Slack</i>	3. Title Analyst, Regulatory Affairs	
4. Typed Name Mandy Slack	5. Date October 24, 2012	

11/13 3/13



**The Scotts Company LLC**

*and Subsidiaries*

October 24, 2012

Tony Kish (PM # 22)  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7505P)  
U.S. Environmental Protection Agency  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202

**SUBJECT: The Scotts Company d/b/a The Ortho Group  
Ortho® Daconil 2787® Multipurpose Fungicide (EPA Reg. No. 239-2522)  
Submission of Label Notification per PR Notice 2008-1 and PR Notice 98-10**

Dear Mr. Kish:

The Scotts Company d/b/a The Ortho Group is submitting a label notification for Ortho® Daconil 2787® Multipurpose Fungicide, 239-2522. In support of this submission, please find enclosed the following materials:

- One (1) fully executed Form 8570-1;
- One (1) "marked up" copy of the master label highlighting the revisions;
- One (1) copy of the revised master label.

The following changes have been made to the enclosed master label:


- Updated the Environmental Hazards section as noted in PR Notice 2008-1.
- Minor labeling additions/corrections per PR Notice 98-10.

If you have any questions concerning this correspondence please contact me by telephone at 937.644.7187 or by email at amanda.slack@scotts.com.

Sincerely,

Mandy Slack  
Analyst, Regulatory Affairs

Enclosures



4/13

EPA Reg. No. 239-2522

NOTE:

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

**ORTHO® DACONIL 2787® Multipurpose Fungicide**

- ABN's: Ortho® Garden Disease Control Concentrate
- Ortho® MAX™ Garden Disease Control Concentrate
- Ortho® Disease B Gon Daconil® Fungicide Concentrate

[] **OUTDOOR USE ONLY**  
 [WATER DROPLET ICON] **WATER BASED**  
**Specially formulated for residential use**

**ACTIVE INGREDIENT:**

Chlorothalonil..... 29.6%

**OTHER INGREDIENTS**..... 70.4%

Total 100.0%

**KEEP OUT OF REACH OF CHILDREN**

**WARNING**

**See back panel booklet for additional precautionary statements**

NET CONTENTS 32 FL OZ (1 QT) 946 mL, 16 FL OZ (1 PT) 473 mL

<b>Product Facts</b>	
<b>STOPS &amp; PREVENTS DISEASES</b>	Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds & other plant diseases
<b>WHERE TO USE/ AMOUNT TO USE</b>	<b>Roses:</b> 1½ tbsp per 4 gal water <b>Other Listed Ornamentals</b> (except conifers & pachysandra): 2 tsp per gal water <b>Vegetables &amp; Fruit Trees:</b> See inside label for list of fruits and vegetables, and dilution rates.
	<b>Questions, Comments or Medical Information [computer icon]</b> call 1-800-225-2883 <a href="http://www.ortho.com">www.ortho.com</a>

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

[Reentry ICON showing adult person & dog] Keep people and pets off treated area until dry.

Do not apply this product in a way that will contact other persons, or pets, either directly or through drift.

**DIRECTIONS:** Readily mixes with water and can be used in an ORTHO® SPRAY-ETTE®, or tank-type sprayer. Fine spray required for proper coverage [; use an ORTHO SPRAY-ETTE® for best results.]  
[ILLUSTRATION - Plant with Brown Patch]

tsp = teaspoon                      tbsp = tablespoon

NOTE:

**Bold, italicized text is information for the reader and is not part of the label.**  
 [Bracketed information is optional text]. Text separated by / denotes and/or options.

EPA Reg. No. 239-2522

VEGETABLES & MELONS

**HOW TO APPLY:** Spray thoroughly to cover all plant surfaces.

VEGETABLES & MELONS DISEASES CONTROLLED	AMOUNT TO USE (for each 200 sq ft)  [Dial-N-Spray applicator illustration] See p. X for Ortho® Dial-n- Spray® directions.	WHEN TO APPLY  Avoid spraying plants during extremely hot & sunny weather. Do not spray wilted plants.	DAYS TO WAIT TO HARVEST
Beans (green) Rust & Gray Mold	2¾ tsp per gal water  4 applications/season	(1) Start spraying during early bloom or at first sign of disease  (2) Apply every 7 days to maintain control.	7
Broccoli, Brussels Sprouts, Cabbage & Cauliflower  • Alternaria Leaf Spot & Downy Mildew  • Ring Spot in Brussels Sprouts (CA only)	• 1½ tsp per gal water  • 2 tsp per gal water  7 applications/season	(1)(a) Start spraying when conditions favor disease; or (b) after transplanting; or (c) when sprouts appear  (2) Apply every 7 days.	0
Carrots Early & Late Blight	2 tsp per gal water  9 applications/season	(1) Start spraying at first sign of disease.  (2) Apply every 7 days to maintain control.	0
Celery  • Early & Late Blight  • Basal Stalk Rot & Pink Rot (suppression)	• 1½ tsp per gal water  • 2¾ tsp per gal water  8 applications/season	(1) Start spraying after transplanting.  (2) Apply every 7 days.	7
Corn (sweet) Helminthosporium Leaf Blight & Rust	1¾ tsp per gal water  6 applications/season	(1) Start spraying when conditions favor disease.  (2) Apply every 7 days to maintain control.	14

6/13

NOTE:

EPA Reg. No. 239-2522

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

VEGETABLES & MELONS DISEASES CONTROLLED	AMOUNT TO USE (for each 200 sq ft)  [Dial-N-Spray applicator illustration] See p. X for Ortho® Dial-n- Spray® directions.	WHEN TO APPLY  Avoid spraying plants during extremely hot & sunny weather. Do not spray wilted plants.	DAYS TO WAIT TO HARVEST
Cucumbers, Melons, Pumpkins & Squash <ul style="list-style-type: none"> <li>• Anthracnose, Downy Mildew &amp; Target Spot</li> <li>• Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Leaf Spot, Powdery Mildew (except southwestern states) &amp; Fruit Rot on Cucumbers (suppression)</li> </ul>	<ul style="list-style-type: none"> <li>• 2 tsp per gal water</li> <li>• 2¾ tsp per gal water</li> </ul> 7 applications/season	(1) Start spraying when plants are in the first true leaf stage or when conditions favor disease.  (2) Apply every 7 days to maintain control.	0
Onion (green and dry bulb); Leek; Shallot & Garlic  Botrytis Leaf Blight (Blast), Purple Blotch & Downy Mildew (suppression)	2¾ tsp per gal water  Onion (green), Leek & Shallot: 3 applications/season  Onion (dry bulb) & Garlic: 7 applications/season	(1) Start spraying at first sign of disease.  (2) Apply every 7 days. However, do not use more than 3 times per season on green onion, leek or shallot.	Onion (green), Leek & Shallot: 14  Onion (dry bulb) & Garlic: 7
Potato Early & Late Blight & Botrytis Vine Rot	1½ tsp per gal water  9 applications/season	(1) Start spraying when plants are 6 inches high or at first sign of disease.  (2) Apply every 5 days to maintain control.	0
Tomato Early & Late Blight, Gray & Septoria Leaf Spots, Gray Leaf Mold, Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot & Botrytis Gray Mold	2½ tsp per gal water  7 applications/season	(1) Start spraying when conditions favor disease.  (2) Apply every 7 days to maintain control.	0

NOTE:

EPA Reg. No. 239-2522

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

**FRUIT TREES**

**HOW TO APPLY: Spray thoroughly to uniformly cover tree canopy (generally sufficient to treat 200 sq. ft.)**

FRUIT TREES DISEASES CONTROLLED	AMOUNT TO USE [Dial-n-Spray applicator illustration] See p. X for Ortho® Dial-n- Spray® directions.	WHEN TO APPLY
Peach, Nectarine & Apricot  <ul style="list-style-type: none"> <li>• Peach Leaf Curl, Coryneum Blight (Shothole)</li> <li>• Scab &amp; Brown Rot Blossom Blight</li> </ul>	3¾ tsp per <b>4-gal</b> water  5 applications/season	Do not apply between fruit formation and harvest.  <ul style="list-style-type: none"> <li>• (1) Spray at leaf fall in late autumn.</li> <li>• (2) Reapply 1 to 2 more times in mid to late winter before buds begin to swell.</li> </ul> <i>Additional directions for Shothole, to prevent fruit infections:</i> (3) Also apply 1 to 2 weeks after petal fall when fruit just begins to form.  <ul style="list-style-type: none"> <li>• (1) Spray when buds begin to show color.</li> <li>• (2) Reapply at full bloom.</li> <li>• (3) If cool, moist weather continues, make a 3rd application at petal fall.</li> <li>• (4) Apply once more when fruit just begins to form.</li> </ul>
Cherry, Plum & Prune <ul style="list-style-type: none"> <li>• Brown Rot Blossom Blight</li> <li>• Cherry Leaf Spot on cherries</li> </ul>	3¾ tsp per <b>4-gal</b> water  5 applications/season	Do not apply between fruit formation and harvest.  (1) Spray when buds begin to show color. (2) Reapply at full bloom. (3) If cool, moist weather continues, make a 3rd application at petal fall.  <i>Additional directions for Cherry Leaf Spot:</i> (4) also apply when fruit just begins to form, and (5) again 7 days after fruit is removed. (6) Apply once more 10 to 14 days later, if tree has a history of leaf spot.
Papaya Alternaria Fruit Spot, Anthracnose & Stem End-Rot	2¾ tsp per <b>4-gal</b> water  3 applications/season	(1) Start spraying when fruit spots appear (April to July). (2) Spray every 14 days until dry weather returns.
Passion Fruit (Hawaii only) Alternaria Fruit & Leaf Spot (Passion Fruit Brown Spot)	1¾ tsp per <b>4-gal</b> water  5 applications/season	(1) Start spraying before spots appear on fruit (April to July). (2) Spray every 14 days until dry weather returns.

8/13

NOTE:

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

EPA Reg. No. 239-2522

ORNAMENTAL PLANTS

**HOW TO APPLY:** Spray thoroughly to cover all plant surfaces (generally sufficient to cover 200 sq. ft.)

**AMOUNT TO USE**

**Roses:** 1½ tbsp per 4-gal water

**Pachysandra:** 3¾ tsp per gal water

**Conifers (Pine, Douglas Fir & Spruce):** 5 tsp per 4-gal water

**Other Ornamental Plants:** 2 tsp per gal water

[Dial-N-Spray applicator illustration] See p. X for Ortho® Dial-n-Spray® directions.

<b>PLANTS DISEASES CONTROLLED</b>	<b>WHEN TO APPLY</b> Apply every 7-14 days (unless otherwise specified) until conditions no longer favor disease. During periods when conditions favor severe disease, generally cloudy or wet weather, use the shorter interval between applications.  <b>SUGGESTED TIMING FOR FIRST APPLICATION</b>
Ash Cercospora, Cercosporidium & Cylindrosporium Leaf Spots	Spring bud break
Azalea & Rhododendron <ul style="list-style-type: none"> <li>• Phytophthora Die-Back</li> <li>• Ovinia Flower Blight</li> </ul> (Discoloration of blooms has been noted on certain varieties when applied during flowering.)	<ul style="list-style-type: none"> <li>• New leaf emergence</li> <li>• Early bloom</li> </ul>
Buckeye (Horsechestnut) Leaf Blotch & Anthracnose	Spring bud break
Carnation <ul style="list-style-type: none"> <li>• Alternaria Leaf Spot</li> <li>• Branch Rot &amp; Botrytis</li> </ul>	<ul style="list-style-type: none"> <li>• Transplant of cuttings</li> <li>• Cool, moist conditions</li> </ul>
Cherry-Laurel Cercospora Leaf Spot	Petal fall
Chrysanthemum (Daisy) Mycosphaerella Ray Blight & Septoria Leaf Spot	Transplant of cuttings
Conifers (Pine, Douglas Fir & Spruce) <ul style="list-style-type: none"> <li>• Needlecast, Rhabdocline &amp; Swiss Needlecast on Douglas Fir</li> <li>• Scirrhia Brown Spot &amp; Scleroderris Canker on Pine</li> <li>• Rhizosphaera Needlecast on Spruce</li> <li>• Lophodermium Needlecast</li> </ul>	<ul style="list-style-type: none"> <li>• (1) Start spraying in spring when new shoots are ½ to 2 inches long.</li> <li>• (2) Spray every 21 days (7 days for seed beds only) until dry weather returns.</li> <li>• North Central &amp; Northeastern States:                (1) Start spraying in mid-July to early August before disease occurs.                (2) Spray every 21 days (7 days for seed beds only) until dry weather returns.</li> </ul>

9/13

NOTE:

EPA Reg. No. 239-2522

**Bold, italicized text is information for the reader and is not part of the label.**  
 [Bracketed information is optional text]. Text separated by / denotes and/or options.

<b>PLANTS DISEASES CONTROLLED</b>	<b>WHEN TO APPLY</b> Apply every 7-14 days (unless otherwise specified) until conditions no longer favor disease. During periods when conditions favor severe disease, generally cloudy or wet weather, use the shorter interval between applications.  <b>SUGGESTED TIMING FOR FIRST APPLICATION</b>
Crabapple Scab, Cedar-apple Rust & Sphaeropsis Leaf Spot	Spring bud break
Dogwood <ul style="list-style-type: none"> <li>• Anthracnose (Discula sp.)</li> <li>• Septoria Leaf Spot</li> </ul>	<ul style="list-style-type: none"> <li>• Spring bud swell</li> <li>• Early bloom</li> </ul>
Dracaena Fusarium Leaf Spot	Pre-transplant
Euonymus Anthracnose	Spring bud break
Flowering Almond, Flowering Quince, Flowering Cherry & Hawthorn <ul style="list-style-type: none"> <li>• Monilinia Blossom/Branch Blight</li> <li>• Rust &amp; Fabraea Leaf Spot</li> </ul>	<ul style="list-style-type: none"> <li>• Early bloom</li> <li>• Pre-bloom</li> </ul>
Geranium <ul style="list-style-type: none"> <li>• Botrytis Flower Blight (Gray Mold)</li> <li>• Rust</li> </ul>	<ul style="list-style-type: none"> <li>• Pre-bloom</li> <li>• Cool, moist conditions</li> </ul>
Gladiolus Curvularia Leaf/Flower Spot & Botrytis Leaf/Flower Spot	Early propagation
Holly Rhizoctonia Web Blight	Warm, moist conditions
Hollyhock Rust	Early seedling stage
Hydrangea (foliage only) Cercospora & Septoria Leaf Spots, & Rust  (Discoloration of blooms has been noted on certain varieties when applied during flowering.)	Early propagation
Iris Botrytis Blossom Blight & Didymellina Leaf Spot	Cool, moist conditions
Leatherleaf Fern Ascochyta & Rhizoctonia Blights, Cercospora & Cylindrocladium Leaf Spots	Spring bud break
Lily Botrytis Gray Mold (foliar phase) & Phytophthora Blight	Pre-bloom
Mountain Laurel Cercospora Leaf Spot	Spring bud break
Oak (red group only) Taphrina Blister, Actinopelte Leaf Spot & Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia) Rust	Spring bud break
Oyster Plant (Rhoeo) Tan Leaf Spot	Early propagation

10/  
13

NOTE:

EPA Reg. No. 239-2522

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional text]. Text separated by / denotes and/or options.

<b>PLANTS DISEASES CONTROLLED</b>	<b>WHEN TO APPLY</b> Apply every 7-14 days (unless otherwise specified) until conditions no longer favor disease. During periods when conditions favor severe disease, generally cloudy or wet weather, use the shorter interval between applications.  <b>SUGGESTED TIMING FOR FIRST APPLICATION</b>
Pachysandra Volutella Leaf Blight	Spring bud break
Parlor Palm (Chamaedorea) Bipolaris Leaf Spot (Helminthosporium)	Cool, moist conditions
Petunia Botrytis Gray Mold (foliar phase), Phytophthora & Botrytis Blights  (Discoloration of blooms has been noted on certain varieties when applied during flowering.)	Pre-bloom
Philodendron Phytophthora Blight & Dactylaria Leaf Spot	Moist conditions
Photinia Fabraea (Entomosporium) Leaf Spot	Spring bud break
Pieris (Andromeda) Phytophthora Die-Back	New leaf emergence
Poplar Marssonina Leaf Spot	Spring bud break
Prayer Plant (Maranta) Helminthosporium Leaf Spot	Early propagation
Privet Cercospora Leaf Spot	Prolonged wet conditions
Pyracantha (Firethorn) Scab	Spring bud break
Rose Black Spot & Botrytis Blight	Spring bud break
Statice Anthracnose, Cercospora & Alternaria Leaf Spots & Botrytis Leaf Blight	Spring bud break
Sycamore (Planetree) Anthracnose	Spring bud break
Syngonium Cephalosporium Leaf Spot	Warm, moist conditions
Viburnum Powdery Mildew	Mid-summer
Zinnia Powdery Mildew	First sign of disease

Combination Spray(s) with Insecticides - on Ornamental Plants

ORTHO® DICONIL 2787 Multi-purpose Fungicide [ORTHO Garden Disease Control] may be mixed with the following insecticides for use on ornamental plants (at the label rates for both products): ORTHO Bug-B-Gon Max Lawn and Garden Insect Killer, ORTHO MAX Malathion Insect Spray Concentrate, or ORTHO® Systemic Insect Control. Apply fungicide on a regular schedule to control disease and add insecticide only when insect control is desired. Do not apply more than three consecutive applications of combination sprays. Do not use combination sprays with Dial-n-Spray or similar devices.

11/  
13

NOTE:

**Bold, italicized text is information for the reader and is not part of the label.**  
[Bracketed information is optional text]. Text separated by / denotes and/or options.

**ORTHO® DIAL-N-SPRAY® DIRECTIONS**

[Dial-N-Spray applicator illustration] TO USE/FOR ORTHO® DACONIL 2787 Multi-purpose Fungicide  
[IN THE ORTHO® DIAL-N-SPRAY®]

1. **Always set dial to 4 oz.**
2. Add product + water to sprayer jar as specified in the tables below.
3. Spray until jar is empty.

Do not store unused spray. Do not pour unused spray back into this Ortho Multi-Purpose Fungicide container.

1 Tablespoon (tbsp) = 3 teaspoons (tsp)    1 fluid ounce (fl oz) = 2 tbsp    8 fl oz = 1 cup

FRUITS, VEGETABLES & ORNAMENTAL PLANTS			
Dial Setting: Always set dial to 4 oz.			
Dilution Rate	Gallons of Spray Produced	Amount of product to add	Amount of water to add
1½ tsp per gal water See Vegetables	1	1½ tsp	7½ tbsp[*]
1¼ tsp per gal water See Vegetables	1	1¼ tsp	7½ tbsp[*]
2 tsp per gal water See Vegetables and Ornamental Plants.	1	2 tsp	7½ tbsp[*]
2½ tsp per gal water See Vegetables	1	2½ tsp	7¼ tbsp[*]
2¾ tsp per gal water See Vegetables.	1	2¾ tsp	7 tbsp[*]
3¼ tsp per gal water See Pachysandra.	1	3¼ tsp	6¾ tbsp[*]
1¾ tsp per <b>4-gal</b> water See Passion Fruit	4	1¾ tsp	15¾ fl oz[**]
2¾ tsp per <b>4-gal</b> water See Papaya	4	2¾ tsp	15½ fl oz[**]
3¾ tsp per <b>4-gal</b> water See Fruit Trees	4	3¾ tsp	15⅓ fl oz[**]
4½ tsp per <b>4-gal</b> water See Roses	4	4½ tsp	15¼ fl oz[**]
5 tsp per <b>4-gal</b> water See Conifers	4	5 tsp	15⅛ fl oz[**]

[Optional Text:

- \*Amount of water will bring volume to 4-oz mark on Dial-n-Spray.
- \*\*Amount of water will bring volume to 16-oz mark on Dial-n-Spray.]

1000179396-01 | In Work - 10/26/2012 by Temp 12/03/2, Melissa McInain | Printed on 10/26/2012 - Print

1000179396\_002\_000\_0101580 prodapp2 MMCLAIN1 2012-10-26T13:52:00 0.945

12/13

NOTE:

EPA Reg. No. 239-2522

**Bold, italicized text is information for the reader and is not part of the label.**  
[Bracketed information is optional text]. Text separated by / denotes and/or options.

**STORAGE AND DISPOSAL [icon]**

**PESTICIDE STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

**PESTICIDE DISPOSAL: If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**CONTAINER DISPOSAL: If empty:** Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING:** Causes substantial but temporary eye injury. Causes skin irritation. Harmful if inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

**Note to User:** Product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation, and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

**FIRST AID**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**Note to Physicians:** Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids. Emergency Information Call 1-800-225-2883.

**ENVIRONMENTAL HAZARDS:**

This pesticide is toxic to aquatic invertebrates and wildlife. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that

13/13

NOTE:

***Bold, italicized text is information for the reader and is not part of the label.***  
[Bracketed information is optional text]. Text separated by / denotes and/or options.

drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

**NOTICE:** To the extent consistent with applicable law, Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

Questions, Comments or Medical Information? Call 1-800-225-2883      www.ortho.com

**The ORTHO Guarantee:** If for any reason you, the consumer, are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

DACONIL 2787® is a Registered Trademarks of GB Biosciences Corporation

2012© The Ortho Group. World rights reserved.

Manufactured for / Sold by:

**The ORTHO Group**  
P.O. Box 190  
Marysville, OH 43040  
EPA Reg. No. 239-2522  
EPA Est. 239-IA-3<sup>1</sup>, 58996-MO-1<sup>A</sup>  
Superscript is first letter of lot number.  
Made in USA

**Optional Marketing Claims**

Makes up to 64 Gallons Diluted Spray [*Pint size only*]  
Makes up to 128 Gallons Diluted Spray [*Quart size only*]

Concentrate  
Controls diseases on: shrubs, vegetables, fruit trees, flowers, and shade trees

CONTAINS DACONIL 2787® FUNGICIDE

CONTROLS DISEASES - Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds, etc. ON:  
Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees.

Controls/Stops & Prevents diseases on vegetables, fruits, roses, flowers, shrubs & trees/ornamentals

Controls/Prevents/Stops garden diseases

Stops & Prevents Garden Disease

Designed for Independent Retailers

For effective control of a broad spectrum of vegetable, fruit, and ornamental plant diseases. Controls:  
Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds, etc. on Vegetables, Fruit Trees, Flowers,  
Shrubs and Shade Trees.

**ORTHO & YOUR ENVIRONMENT**

Your home and yard are places for family and pets to enjoy. That's why Ortho® products are designed with care to provide effective solutions to garden disease problems. For best results and a healthy environment, please follow instructions for appropriate usage, storage and disposal.

[OPEN Resealable Label for Directions & Precautions] [PRESS TO RESEAL]

1000179396-01 | In Work - 10/26/2012 by Temp 12/03/2, Melissa McLain | Printed on 10/26/2012 - Print

1000179396\_002\_000\_0101580 prodapp2 MMCLAIN1 2012-10-26T13:52:00 0.945